

BREEDING BIRD CENSUS

Last updated November 2009

Chris & Lynda Huxley

Since 2003, we have been carrying out a census of the birds present in the breeding season in three areas around the shores of Lough Carra: the Doon peninsula, the Kilkeeran peninsula and the Partry House Estate. These three locations were chosen primarily to include examples of major terrestrial habitat types: woodland, fen, grassland, scrub and hedgerow, as well as the lakeshore and aquatic habitats (such as reedbed and open water).

The main objective of this regular census is:

To monitor the populations of bird species breeding, or possibly breeding, around Lough Carra.

In addition, the census provides a baseline inventory of bird species occurring around the lake in April/May as well as a comparative indication of bird diversity in these areas.

The methodology used is based on a standard technique for such monitoring work when only a relative estimate of abundance (a population “index”) is wanted: each location is censused twice in the breeding season, usually in late April and late May. A fixed transect route is walked by two observers who record every bird that is positively identified by sight or by song or call within approximately 100 metres of the route. Each transect is covered at approximately the same rate and weather conditions (dry and at least partially sunny) in the early morning between 06.30 and 09.30. The same two observers have carried out this work each year so far, but the method is such that results from other experienced observers should be broadly comparable.

For further background on such methods, see Bibby et al. 1998.

The Transects

1. At the Doon peninsula, the transect route starts at the small “car park” area (grid ref. M 172760), follows the track into and through the woodland to the ancient fortified wall, and then follows the path around the peninsula until returning to the fortified wall, where the transect ends. The time taken for this route is approximately 1 hour 15 minutes. The main habitats include grassland, mature deciduous woodland and scrub as well as the lakeshore and the lake itself.
2. At the Kilkeeran peninsula, the transect starts at the far end of the causeway (grid ref. M 171712) and follows the lakeshore in an anticlockwise direction, ending at the starting point. The time taken for this is approximately 1 hour 15 minutes and the main habitats include grassland, fen, scrub, hedgerow, a small area of woodland and some reedbed, as well as the lakeshore, a couple of small islands and the lake itself.

3. On the Partry House Estate, the transect starts at Partry House (grid ref. M 166695), follows a northerly direction across farmland to the woodland edge (grid ref. M 166700), turns easterly along the woodland edge to M 168700, crosses the fen to the northern tip of Creggaun wood (M 171699), follows the path through Creggaun wood, across the fen via the “stepping stones” and then follows the lakeshore around the eastern side of Moynish to the southernmost point (M 169688), after which the main path north through the woodland is taken, to join the “famine causeway” at M 169695. The route follows the causeway and then the lakeshore as far as the Keel River at M 163682, where the transect ends. This route takes around three hours to complete and in some conditions the transect is completed in two parts over two days in order to avoid continuing after about 09.30. The habitat types covered include deciduous woodland, mixed coniferous/deciduous woodland, scrub, grassland, hedgerow, fen, reedbed and the lake.

Thus, each year a total of 11 hours of transect counts is recorded.

The Results

The data collected through these surveys are presented below. However, there are several important points that must be stressed when examining and interpreting the figures:

- Two species are specifically excluded: the Black-headed gull and Common gull, as they both breed colonially in considerable numbers on islands at the Kilkeeran peninsula.
- The count data for **both** the April and May transects for each location are combined. For this reason (and because the method is, in any case, not suitable), the figures should not be used to estimate the number of birds or pairs of a species present in the area. Thus, the figures should be used only as an **index** of abundance. For example, if 12 Great crested grebes are recorded at Kilkeeran in one year, this is the total recorded in **both** census walks combined. It does **not** indicate that there were six pairs, nor that there were three pairs (i.e. 6 birds in April and the same 6 in May). Its value lies in the fact that, if the same methodology is used each year, any change in this number may be interpreted as a change in overall abundance of this species in the areas surveyed.
- The data for those species recorded in only very small numbers must be interpreted with great caution. The methodology is such that the presence of a species that occurs in only very small numbers may well be missed in one year, but recorded in another. For example, Jays occur regularly in two of the three locations, but may well be missed in a census if the timing is such that they are in another part of the woodland during the brief census period.
- The likelihood of recording large species or those with loud vocalisations is greater than for small and/or “quiet” species. This variation is increased in those habitats or areas where observation distances are limited (especially in mature woodland). Great care should be taken, therefore, when comparing relative abundance of different species.

- In 2007, the census was incomplete as factors beyond our control prevented the Partry transect from being carried out. This does not affect the results for Doon and Kilkeeran, but prevents compilation of totals for 2007.
- In all years and on all transects, species may be recorded that do not breed in the area. Most of these will be passage migrants or winter visitors that have yet to leave. Golden plover, for example, are sometimes recorded in considerable numbers at Kilkeeran, but they do not breed there. Care must be taken therefore, in interpreting the data for such species.

REFERENCES

Bibby, C., M. Jones and S. Marsden. 1998. Expedition Field Techniques: Bird Surveys. Royal Geographical Society, London.

Chris & Lynda Huxley
Updated November 2009

Breeding Bird Census by Site

Species	2003	2004	2005	2006	2007	2008	2009
	Doon	Doon	Doon	Doon	Doon	Doon	Doon
Little grebe							
Great-crested grebe				2			
Cormorant		1			1		
Grey heron	1						
Mute swan							
Barnacle goose							
Wigeon							
Gadwall							
Teal							
Mallard	2	3	8			3	10
Shoveler							
Tufted duck						3	
Red-breasted merganser	2						
Sparrowhawk		1					
Kestrel							
Peregrine							
Pheasant							
Water rail							
Moorhen							
Coot							
Oystercatcher							
Ringed plover							
Golden plover							
Lapwing							
Dunlin							
Snipe							
Black-tailed godwit							
Whimbrel							
Curlew						7	
Redshank							
Common sandpiper							
Common tern							1
Woodpigeon	9	7	2	5	8	20	9
Collared dove							
Cuckoo	1						
Swift						1	
Kingfisher							
Skylark							
Sand martin	1					3	
Swallow	1						1
House martin							
Meadow pipit							
Grey wagtail							
Pied wagtail							
Wren	13	10	12	22	12	16	8
Dunnock	2			3			
Robin	11	21	12	18	15	9	15
Stonechat							
Wheatear							
Blackbird	11	14	13	8	11	12	9
Song thrush	10	20	3	5	5	13	5
Mistle thrush	1						
Grasshopper warbler							
Sedge warbler							

Whitethroat							
Blackcap	10	4	4	7	9	10	15
Chiffchaff	2	5	2	2	3	1	1
Willow warbler	33	23	24	22	28	15	30
Goldcrest	2	7	4	6	2	2	3
Long-tailed tit		7	2		4		
Coal tit	2	5	4	2	1	1	
Blue tit		5	7	10	8	13	13
Great tit	5	3	8	3	13	5	5
Treecreeper	2		1				
Jay		1	1	2	1		1
Magpie	1	1					
Jackdaw						1	2
Rook	4	1				2	
Hooded crow	2	8	1	3	2	4	1
Raven	2						
Starling							
House sparrow							
Chaffinch	18	12	13	16	14	17	16
Greenfinch							
Goldfinch							
Linnet							
Redpoll							
Bullfinch	2	2	6	2	7	4	2
Reedbunting	7	7	4	12	3	4	1
TOTAL	157	168	131	150	147	166	148
Number of Species	27	22	20	19	19	23	20
	19-Apr	18-Apr	16-Apr	19-Apr	24-Apr	28-Apr	25-Apr
	29-May	14-May	14-May	21-May	27-May	27-May	29-May

Breeding Bird Census by Site

Species	2003	2004	2005	2006	2007	2008	2009
	Kilkeeran	Kilkeeran	Kilkeeran	Kilkeeran	Kilkeeran	Kilkeeran	Kilkeeran
Little grebe					1	2	
Great-crested grebe	9	6	11	10	20	25	17
Cormorant	1						1
Grey heron		1					
Mute swan	4		4	2	4	4	1
Barnacle goose			1				
Wigeon					1		
Gadwall	5			1		1	2
Teal							
Mallard	19	7	6	10	10	30	28
Shoveler					2		
Tufted duck	21	8	21	23	73	68	50
Red-breasted merganser	7						
Sparrowhawk							
Kestrel							
Peregrine							
Pheasant						1	1
Water rail							
Moorhen				2		2	2
Coot	3		8	3	6	4	2
Oystercatcher			1				
Ringed plover	8	9	5	1			
Golden plover	87	33					
Lapwing	23	8	5	4	4	4	2
Dunlin			2				
Snipe	8	4	5	4	1	2	4
Black-tailed godwit							
Whimbrel			17			5	7
Curlew					8		
Redshank	10	3	5	2	5	7	2
Common sandpiper		1	1		1		5
Common tern				2		2	2
Woodpigeon	6	1	4	2	6	2	8
Collared dove							
Cuckoo	1				1		
Swift							
Kingfisher							
Skylark	3	3			2	1	1
Sand martin	2	4					
Swallow	3			1	1		
House martin							
Meadow pipit	6	3	4	10	10	9	6
Grey wagtail							
Pied wagtail							
Wren	11	20	16	11	14	18	7
Dunnock	3	3	3	2	1		
Robin	5	2	7	13	12	14	14
Stonechat							
Wheatear					1		
Blackbird	20	11	25	19	25	25	23
Song thrush	8	6	8	6	8	7	4
Mistle thrush							
Grasshopper warbler							
Sedge warbler		1		3		4	2

Whitethroat	1					2	1
Blackcap						2	
Chiffchaff							
Willow warbler	11	14	15	19	12	8	8
Goldcrest				1		3	
Long-tailed tit			1				
Coal tit						1	
Blue tit	3	2	5	7		1	
Great tit				3	1		1
Treecreeper							
Jay							
Magpie	2						
Jackdaw	2						
Rook							
Hooded crow			2	2	5	1	
Raven							
Starling					6	10	
House sparrow							
Chaffinch	15	11	12	7	5	11	9
Greenfinch		2		1		1	
Goldfinch							3
Linnet	2	10	9		4	4	
Redpoll				2			
Bullfinch		3	2	2			3
Reedbunting	15	23	22	24	9	25	20
TOTAL	324	199	227	199	259	306	236
Number of Species	32	27	30	31	31	34	30
	21-Apr	24-Apr	29-Apr	23-Apr	26-Apr	28-Apr	2-May
	28-May	18-May	15-May	30-May	29-May	27-May	30-May

Breeding Bird Census by Site

Species	2003	2004	2005	2006	2007	2008	2009
	Partry	Partry	Partry	Partry	Partry	Partry	Partry
Little grebe						2	
Great-crested grebe							
Cormorant							2
Grey heron			1			2	
Mute swan						4	
Barnacle goose							
Wigeon							
Gadwall							
Teal							
Mallard	1	17	5	14		28	15
Shoveler							
Tufted duck		4		4		2	7
Red-breasted merganser							
Sparrowhawk						1	
Kestrel						1	1
Peregrine		1					
Pheasant		1	1			2	1
Water rail	1	1	1				1
Moorhen		1	3			6	2
Coot	4	1	3			4	2
Oystercatcher							
Ringed plover	3	2	2	2		4	
Golden plover							
Lapwing		3	2				
Dunlin							
Snipe	1	1		2		1	
Black-tailed godwit			6				
Whimbrel							
Curlew							
Redshank	1	1					
Common sandpiper	4	2	3	5		4	2
Common tern				3			1
Woodpigeon	42	30	24	22		18	19
Collared dove		1	1				
Cuckoo	1		1				2
Swift							
Kingfisher							
Skylark							
Sand martin		1					
Swallow	3	11	6	4		17	13
House martin							
Meadow pipit				1			
Grey wagtail		1		1			
Pied wagtail		4				5	2
Wren	30	32	36	28		32	21
Dunnock	2	1		1			
Robin	40	35	38	34		23	25
Stonechat							
Wheatear							
Blackbird	33	39	30	26		33	24
Song thrush	11	26	25	17		22	15

Mistle thrush	1	5	8	16		6	1
Grasshopper warbler	1	1				1	
Sedge warbler	3	5	3	3		9	3
Whitethroat				1			
Blackcap	8	14	5	8		13	4
Chiffchaff	5	6	3	5		5	3
Willow warbler	32	32	31	27		23	27
Goldcrest	8	15	16	14		16	9
Spotted flycatcher						7	
Long-tailed tit	2		3			2	
Coal tit	12	10	7	3		3	5
Blue tit	14	13	26	20		15	6
Great tit	10	8	13	8		6	6
Treecreeper	2		4	2		4	
Jay				2			
Magpie			2			1	3
Jackdaw	18	1		8		4	
Rook						1	
Hooded crow	4		3	3		8	2
Raven							1
Starling	10	18	24	7		27	11
House sparrow							
Chaffinch	31	25	45	27		27	20
Greenfinch							
Goldfinch			1			2	1
Linnet							
Redpoll	2					1	
Bullfinch		5		6		6	4
Reedbunting	11	3	9	7		8	5
TOTAL	351	377	391	331	-	406	266
Number of Species	33	35	34	33	-	43	35
	22-Apr	25-Apr	24-Apr	20-22-Apr	no count	27-Apr	26-Apr
	31-May	15-May	22-26 May	28-May	no count	1-Jun	24-May

Breeding Bird Census Totals

Species	2003	2004	2005	2006	2007	2008	2009
	Total	Total	Total	Total	Total	Total	Total
Little grebe						4	
Great-crested grebe	9	6	11	12		25	17
Cormorant	1	1					3
Grey heron	1	1	1			2	
Mute swan	4		4	2		8	1
Barnacle goose			1				
Wigeon							
Gadwall	5			1		1	2
Teal							
Mallard	22	27	19	24		61	53
Shoveler							
Tufted duck	21	12	21	27		73	57
Red-breasted merganser	9						
Sparrowhawk		1				1	
Kestrel						1	1
Peregrine		1					
Pheasant		1	1			3	2
Water rail	1	1	1				1
Moorhen		1	3	2		8	4
Coot	7	1	11	3		8	4
Oystercatcher			1				
Ringed plover	11	11	7	3		4	
Golden plover	87	33					
Lapwing	23	11	7	4		4	2
Dunlin			2				
Snipe	9	5	5	6		3	4
Black-tailed godwit			6				
Whimbrel			17			5	7
Curlew						7	
Redshank	11	4	5	2		7	2
Common sandpiper	4	3	4	5		4	7
Common tern				5		2	4
Woodpigeon	57	38	30	29		40	36
Collared dove		1	1				
Cuckoo	3		1				2
Swift						1	
Kingfisher							
Skylark	3	3				1	1
Sand martin	3	5				3	
Swallow	7	11	6	5		17	14
House martin							
Meadow pipit	6	3	4	11		9	6
Grey wagtail		1		1			
Pied wagtail		4				5	2
Wren	54	62	64	61		66	36
Dunnock	7	4	3	6			
Robin	56	58	57	65		46	54
Stonechat							
Wheatear							
Blackbird	64	64	68	53		70	56
Song thrush	29	52	36	28		42	24
Mistle thrush	2	5	8	16		6	1
Grasshopper warbler	1	1				1	

Sedge warbler	3	6	3	6		13	5
Whitethroat	1			1		2	1
Blackcap	18	18	9	15		25	19
Chiffchaff	7	11	5	7		6	4
Willow warbler	76	69	70	68		46	65
Goldcrest	10	22	20	21		21	12
Spotted flycatcher						7	
Long-tailed tit	2	7	6			2	
Coal tit	14	15	11	5		5	5
Blue tit	17	20	38	37		29	19
Great tit	15	11	21	14		11	12
Treecreeper	4		5	2		4	
Jay		1	1	4			1
Magpie	3	1	2			1	3
Jackdaw	20	1		8		5	2
Rook	4	1				3	
Hooded crow	6	8	6	8		13	3
Raven	2						1
Starling	10	18	24	7		37	11
House sparrow							
Chaffinch	64	48	70	50		55	45
Greenfinch		2		1		1	
Goldfinch			1			2	4
Linnet	2	10	9			4	
Redpoll	2			2		1	
Bullfinch	2	10	8	10		10	9
Reedbunting	33	33	35	43		37	26
TOTALS	832	744	749	680	*	878	650
Number of Species	51	52	49	42	*	57	47

* 2007 Totals unavailable - census incomplete